

Random Schrödinger operators on long boxes, noise explosion and the GOE

Benedek Valkó^{*} Bálint Virág[†]

July 26, 2011

Abstract

It is conjectured that the eigenvalues of random Schrödinger operators at the localization transition in dimensions $d \geq 2$ behave like the eigenvalues of the Gaussian Orthogonal Ensemble (GOE). We study a the eigenvalues of long boxes in dimension $d = 2$ for low disorder. We deduce a stochastic differential equation representation for the limiting process. We show that in dimension $d = 2$ there are sequences of boxes so that the eigenvalues in low disorder converge to Sine₁, the limiting eigenvalue process of the GOE.

1 Introduction

When Wigner (1957) introduced random matrices to model large atomic nuclei, his main goal was to find a simple mathematical model that shows repulsion between eigenvalues as observed in the data. The Gaussian orthogonal ensemble (GOE) has since been a remarkable success in the physics literature: several important complex systems are predicted to have eigenvalue repulsion akin to the GOE, most notably the Laplacians of many planar domains, see Bohigas et al. (1984).

Given the all the non-rigorous theories that predict GOE behavior, as well as a lot of numerical evidence, it is surprising that there are hardly any mathematically rigorous results in this direction. Most objects with rigorously known bulk GOE behavior are themselves

^{*}Department of Mathematics, University of Wisconsin - Madison, WI 53705, USA. valko@math.wisc.edu

[†]Department of Mathematics, University of Toronto, ON M5S 2E4, Canada. balint@math.toronto.edu

random matrices constructed similarly to the Gaussian orthogonal ensemble, see Tao and Vu (2009) and Erdős et al. (2009a) for recent breakthrough results in this direction.

The goal of this paper is to show that random Schrödinger operators on long boxes in $\mathbb{Z}^2$ behave like the Gaussian orthogonal ensemble. More precisely, we will show

Theorem 1. *There exists a sequence of weighted boxes on $\mathbb{Z}^2$ with diameter converging to ∞ so that the rescaled eigenvalue process of the adjacency matrix plus suitable diagonal noise converges to the Sine_1 point process.*

Recall that the Sine_1 point process is the limit of the eigenvalues at the bulk of the spectrum of the Gaussian orthogonal ensemble, see Mehta (2004). The process-level convergence described here implies convergence of eigenvalue gap sizes and all similar local statistics.

The concept of localization for random Schrödinger operators was introduced by Anderson (1958). As more and more diagonal noise is added to the Laplacian of a large box, the eigenvectors change from being spread out over the entire box to localized on smaller regions. It is believed (see, for example, Altshuler and Shklovski (1986), Efetov (1997)) that near this transition the eigenvalue process is GOE-like, and, as the noise increases, it is approximately Poisson. This transition from GOE to Poisson can be analyzed in the long-box case. We have

Theorem 2 (Eigenvalue process transition). *Consider boxes in $\mathbb{Z}^2$ with fixed base $\mathbb{Z}_m$. Consider the process of eigenvalues of*

$$n\left(\frac{1}{2}\mathbb{Z}_m \times \mathbb{Z}_n + \frac{\sigma}{\sqrt{n}}V - I\lambda^*\right),$$

where V are diagonal matrices with independent, mean zero, variance 1 entries with bounded third moment. Then

- *For almost all $\lambda^* \in [0, 1]$, for appropriate subsequences, this process has a limit Λ_σ which depends on m, λ^* .*
- *As $\sigma \rightarrow 0$, the process Λ_σ converges to the (randomly shifted) eigenvalue process of a random matrix with independent Gaussian entries.*
- *For every σ , the process Λ_σ is the zero process of the determinant of a $m+1$ -dimensional matrix-valued analytic function described by a stochastic differential equation.*

Remark 3. To illustrate the GOE-to-Poisson transition, it is possible to show that

- *as $\sigma \rightarrow \infty$ the process Λ_σ converges to the Poisson point process.*

We plan to do this in a subsequent article.

The proofs of both theorems are based on the analysis of transfer matrices. The more precise statements can be found in Proposition 4, Corollary 6, Proposition 9, and Corollary 11.

The noise term in the transfer matrix evolution in this scaling regime for block Jacobi matrices has been studied in Römer and Schulz-Baldes (2010) using a language different from SDEs. The first arxiv version of the present paper was followed by the preprint of the paper Bachmann and De Roeck (2010), who, in independent work, also study SDE limits of transfer matrices. Their starting point is the so-called DMPK theory in the physics literature, which is essentially the study of diffusive limits of quasi-one-dimensional random Schrödinger operators from a slightly different point of view. We refer the reader to Bachmann and De Roeck (2010) for a discussion of this theory. One of the novelties of our approach is that it allows for studying the dependence on the eigenvalue λ , which in turn allows us to deduce the scaling limit of the spectrum, the main focus here.

Noise explosion. The central reason for the appearance of random matrices is noise explosion. For a simple case of this, consider

$$B_{\eta,n}(t) = \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor} e^{i\eta k} X_k,$$

where the X_k are independent random variables, with, say, standard normal distribution. If $\eta = 0 \pmod{2\pi}$, then the sequence of functions $B_{\eta,n}$ converges in law to real Brownian motion. Otherwise, $B_{\eta,n}$ converges to complex Brownian motion. Thus a one-dimensional noise process X_k gives rise to a two real-dimensional Brownian motion in the presence of oscillatory terms. Moreover, if η, ν are linearly independent over the integers, then $B_{\eta,n}, B_{\nu,n}$ converge jointly to two independent Brownian motions.

Now we change the setting so that the X_k are $m \times m$ diagonal matrices with independent standard normal entries. Let U be a unitary matrix with eigenangles that are linearly independent over the integers, and also assume that the absolute squares of the eigenvectors of U are not orthogonal to each other. Now consider the sum

$$B_n(t) = \frac{1}{\sqrt{n}} \sum_{k=0}^{\lfloor nt \rfloor} U^{-k} X_k U^k.$$

A simple computation shows that $B_n(t)$ converges to a Hermitian matrix-valued Brownian motion process which is m^2 -real dimensional. In such oscillatory sums, noise that was originally restricted to the diagonal explodes into all possible directions, and changes dimension

from m to m^2 . This phenomenon, which we call **noise explosion**, plays a central role in the proofs below.

The method of analysis in this paper is an extension of the 1-dimensional case introduced in a different setting in Valkó and Virág (2009) and further refined and simplified in Kritchevski, Valkó, and Virág (2011). The latter paper studies the most natural 1-dimensional random Schrödinger operator at the localization transition.

The current paper does not give any bounds on the aspect ratio of the long boxes m/n . This would require rates of convergence estimates for diffusion approximation.

Problem 1. Show that the result in this paper holds for $n = O(m^q)$ for some $q \geq 1$. What is the smallest possible value of q for which the results hold? Is it $q = 1$?

Problem 2. Extend the results to long boxes of higher dimension. The eigenvalue structure of the base here is more complicated, see Remark 12.

Problem 3. Eliminate the $r \rightarrow 0$ condition in Proposition 9. This would require more general versions of the results in Erdős et al. (2010).

2 Description of the model and notation

Let $\mathbb{Z}_n$ denote the graph of a path of length n with vertices $1, 2, \dots, n$. We will use the same variables for a (weighted) graph and its adjacency matrix, so $\mathbb{Z}_n$ will also denote the $n \times n$ matrix with entries $(\mathbb{Z}_n)_{i,j} = 1_{\{|i-j|=1\}}$. Let $\times$ denote the cartesian product of weighted graphs (or matrices), namely

$$(A \times B)_{(i,i'),(j,j')} = \mathbf{1}_{i'=j'} A_{i,j} + \mathbf{1}_{i=j} B_{i',j'}$$

For a positive real r and weighted graph (or matrix) G we will denote by rG the graph where the weights are all multiplied by r (which is the same as multiplying the matrix G by r).

Boxes. The boxes we consider are $[r(\mathbb{Z}_m)] \times \mathbb{Z}_n$ for some positive real r , where n is typically much larger than all the m_i . This is the adjacency matrix of a weighted graph on the box $\mathbb{Z}_m \times \mathbb{Z}_n$ where the edges in the first direction have weight r , and the edges in the second direction have weight 1.

The adjacency matrix corresponds to Dirichlet boundary conditions. For a probabilistic interpretation, the matrix

$$r\mathbb{Z}_m \times \mathbb{Z}_n - 2(r+1)I_{m \times n}$$

is the transition probability matrix of the continuous time random walk on the box killed when it leaves; the jump rates are r and 1 in the two different kinds of directions.

General slabs. Some of our results will apply to $rG \times \mathbb{Z}_n$ where G is a general symmetric matrix.

Chaoticity. For the noise explosion phenomenon to work, we need to consider conjugation by special angles. Consider points (angles) $\mathcal{X} = \{x_1, \dots, x_m\}$ on the unit circle $\mathbb{R}/2\pi$. We will be interested in the set

$$\mathcal{A} := \{x_{i_1} + x_{i_2} + x_{i_3} + x_{i_4}\} \cup \{x_{i_1} + x_{i_2} - x_{j_1} - x_{j_2} : i_k \neq j_\ell\} \cup \{x_{i_1} + x_{i_2} + x_{i_3} - x_{j_1}\}$$

where the indices run through $1, \dots, m$ and the operations are meant mod 2π . The **chaoticity** of $\mathcal{X}$ is defined as the distance of the set $\mathcal{A}$ from 0

$$\text{cha}(\mathcal{X}) = \text{dist}(0, \mathcal{A}) \tag{1}$$

We call the x_i **chaotic** if they have nonzero chaoticity.

In particular, if the $\mathcal{X}$ are chaotic then the x_i 's are distinct. Moreover, if x_i 's are linearly independent over the integers then they are chaotic. A condition very similar to chaoticity appears as the main assumption in Schulz-Baldes (2004).

The **noise explosion** described in the introduction will rely on the following quantities. Let $G = ODO^{-1}$ be the diagonalization of G , so that O is orthogonal and D is diagonal. Let $O_i = O_{i,\cdot}$ denote the i th row of O . Let $|O_i|^2$ denote the vector formed by the squares of the entries of O_i .

When G is the adjacency matrix of the discrete interval, then the eigenvectors are of the form

$$O_{jk} = \sqrt{2/(m+1)} \sin(\pi jk/(m+1)),$$

and it is easy to check that

$$(m+1)\langle |O_i|^2, |O_j|^2 \rangle = \begin{cases} 3/2 & \text{for } i = j, \\ 1 & \text{for } i \neq j. \end{cases} \tag{2}$$

3 The regularized transfer matrix evolution

For this section, consider the operator $G \times \mathbb{Z}_n$ where G is a symmetric matrix of dimension $m \times m$.

Suppose that we want to solve the equation $Mu = \lambda u$ where $M = G \times \mathbb{Z}_n + V$ with a diagonal matrix V . We will do this by solving the system of linear equations recursively,

slice by slice (where a slice is a copy of G , i.e. $G \times \{i\}$). Let u be a function from the vertices of $G \times \mathbb{Z}_n$ to $\mathbb{R}$, and let $u_i(j) = u(j, i)$ so that u_i is a vector indexed by G . Denote by V_i the diagonal matrix which is the restriction of V to the indices $G \times \{i\}$. Then the eigenvalue equation for entries with the same first coordinate $2 \leq i \leq n-1$ reads

$$\lambda u_i = (G + V_i)u_i + u_{i-1} + u_{i+1} \quad (3)$$

and if we set $u_0 \equiv u_{n+1} \equiv 0$, then (3) holds for $i = 1$ and n as well. It is not hard to check that this system of n equations is then equivalent to $Mu = \lambda u$. Note that (3) can be written in terms of block transfer matrices of dimension $2m$ as follows:

$$\begin{bmatrix} u_{k+1} \\ u_k \end{bmatrix} = \begin{bmatrix} \lambda I - G - V_k & -I \\ I & 0 \end{bmatrix} \begin{bmatrix} u_k \\ u_{k-1} \end{bmatrix}.$$

Denoting the above matrix by T_k , we can now characterize the eigenvalues of M as

$$\{\lambda : \exists u_1, u_n \text{ such that } T_n \cdots T_2 T_1 \begin{bmatrix} u_1 \\ 0 \end{bmatrix} \parallel \begin{bmatrix} 0 \\ u_n \end{bmatrix}\} = \{\lambda : \det((T_n \cdots T_2 T_1)_{(1,1)})\}. \quad (4)$$

Here the subscript $(1,1)$ refers to the top left $m \times m$ submatrix, and both representations are equivalent to this submatrix having some eigenvector u_1 with eigenvalue 0. So we may write

$$\text{spec}(M) = \text{zeros}_\lambda [\det((T_n \cdots T_2 T_1)_{(1,1)})] \quad (5)$$

where $\text{spec}(\cdot)$ will refer to the eigenvalue counting measure and zeros_λ refers to the zero counting measure of a real-analytic function of λ . It is not hard to check that the equality also holds in the presence of multiple zeros.

We now introduce the unperturbed version of the transfer matrix

$$T_* = \begin{bmatrix} \lambda_* I - G & -I \\ I & 0 \end{bmatrix}.$$

where λ_* is some fixed reference point. Let $G = ODO^T$ be the diagonalization of G with O orthogonal and $D = \text{diag}(d_1, \dots, d_m)$ diagonal. We will use the shorthand notation $X^Y = Y^{-1}XY$ which means $G = D^{O^T}$.

We may change basis to study products of

$$T_k^{O \otimes I_2} = \begin{bmatrix} \lambda I - D - V_k^O & -I \\ I & 0 \end{bmatrix}, \quad O \otimes I_2 = \begin{bmatrix} O & 0 \\ 0 & O \end{bmatrix}.$$

and note that

$$T_*^{O \otimes I_2} = \begin{bmatrix} \lambda_* I - D & -I \\ I & 0 \end{bmatrix}$$

This matrix can be completely diagonalized as it is just the permutation of a block diagonal matrix with 2×2 blocks. (We have learned that such a change of basis has been considered for a slightly different problem by Schulz-Baldes (2004).) Let

$$z_j = (\lambda_* - d_j)/2 + i\sqrt{1 - ((\lambda_* - d_j)/2)^2}$$

and $Z = \text{diag}(z_1, \dots, z_m)$, We assume that $|\lambda_* - d_j| \in (0, 2)$ for all j which means that z_j is a unit length complex number in the upper half plane.

Then we have

$$T_*^{(O \otimes I_2)Q} = \begin{bmatrix} \bar{Z} & 0 \\ 0 & Z \end{bmatrix}.$$

where

$$S = \begin{bmatrix} \text{diag} |\bar{z}_j - z_j|^{-1/2} & 0 \\ 0 & \text{diag} |\bar{z}_j - z_j|^{-1/2} \end{bmatrix}, \quad Q = \begin{bmatrix} \bar{Z} & Z \\ I & I \end{bmatrix} S \quad (6)$$

and

$$Q^{-1} = iS \begin{bmatrix} I & -Z \\ -I & \bar{Z} \end{bmatrix}$$

Then the matrix

$$X_k = (T_*^{-k} T_k \cdots T_2 T_1)^{(O \otimes I_2)Q},$$

is deterministic matrix factors away from the transfer matrix evolution $T_k \cdots T_1$. However, it has the advantage that, unlike the product $T_k \cdots T_1$, it changes slowly as k varies. This can be seen from its evolution. From the definition we have

$$X_k = (T_*^{-k} T_k T_*^{-1} T_*^k)^{(O \otimes I_2)Q} X_{k-1}, \quad (7)$$

Note that in our case T_k will be a small perturbation of T_* .

We call $(X_k, 0 \leq k \leq n)$ the **regularized transfer matrix evolution**, regularized at λ_* . In the next section, we will show that this evolution has a stochastic differential equation limit. But first let us check how to read off the eigenvalues of the operator M from X_n . By (5) the eigenvalues of M are given by

$$\begin{aligned} \left\{ \lambda : \det \left(((T_*^{(O \otimes I_2)Q})^n X_n)^{Q^{-1}(O^{-1} \otimes I_2)} \right)_{(1,1)} = 0 \right\} \\ = \left\{ \lambda : \det \left(((T_*^{(O \otimes I_2)Q})^n X_n)^{Q^{-1}} \right)_{(1,1)} = 0 \right\} \end{aligned}$$

Note that $(T_*^{(O \otimes I_2)Q})^n X_n$ is a matrix of the form

$$\begin{bmatrix} \bar{Z}^n X_{11} & \bar{Z}^n X_{12} \\ Z^n X_{21} & Z^n X_{22} \end{bmatrix},$$

and so the $(1, 1)$ block of $Q(T_*^{(O \otimes I_2)Q})^n X_n Q^{-1}$ is given by

$$iS_{11}(\bar{Z}^{n+1} X_{11} - \bar{Z}^{n+1} X_{12} + Z^{n+1} X_{21} - Z^{n+1} X_{22})S_{11} = iS_{11}(\bar{Z}^{n+1}, Z^{n+1})X_n(I, -I)^T S_{11}$$

Since S_{11} is nonsingular, have

$$\begin{aligned} \text{spec}(M) &= \text{zeros}_\lambda [\det((\bar{Z}^{n+1}, Z^{n+1})X_n(I, -I)^T)] \\ &= \text{zeros}_\lambda [\det \Im(\bar{Z}^{n+1}((X_n)_{11} - (X_n)_{12}))]. \end{aligned} \quad (8)$$

the second equality follows from the fact that the determinant is zero only for real λ (as these are eigenvalues of a symmetric matrix M), and for real λ , we have $X_{22} = \bar{X}_{11}$ and $X_{21} = \bar{X}_{12}$. This will be shown in the proof of Proposition 4 in the next section.

4 The limiting of the transfer matrix evolution

The goal of this section is to show that the regularized transfer matrix evolution introduced in the previous section has a stochastic differential equation limit. It will follow that the eigenvalues of the corresponding operator converge in distribution to an explicitly constructible limit.

We first state the scaling limit of the evolution of transfer matrices.

Proposition 4 (Limiting evolution of transfer matrices). *Fix $r, \sigma > 0$. Consider the operator $M_n = rG \times Z_n + n^{-1/2}\sigma V$, where $V = V^{(n)}$ is a diagonal matrix with independent random entries of mean 0, variance 1 and uniformly bounded third moments. Denote the eigenvalues of G by $d_1, \dots, d_m$ and assume that $0 < |\lambda_* - d_j| < 2$ for all j and that the angles $\arccos((\lambda_* - rd_j)/2)$, $j = 1, \dots, m$ are chaotic. Consider the regularized transfer matrix evolution corresponding to λ_* : $(X_k^{(n)}, 0 \leq k \leq n)$.*

Then for any finite $\Lambda \subset \mathbb{C}$ we have convergence in distribution as $n \rightarrow \infty$:

$$(X_{[nt]}^{(n)}(\lambda^* + \lambda/n), t \in [0, 1], \lambda \in \Lambda) \Rightarrow (Y_t(\lambda), t \in [0, 1], \lambda \in \Lambda)$$

where Y is the solution of the SDE

$$dY = S^2 \begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix} i\lambda dt Y + i\sigma S \begin{bmatrix} dA & dB \\ -d\bar{B} & -d\bar{A} \end{bmatrix} SY, \quad Y_0 = I. \quad (9)$$

and S is defined in (6). Moreover, for any fixed t , we have convergence in distribution of $X_{[nt]}^{(n)}(\cdot) \rightarrow Y_t(\cdot)$ with respect to the uniform-compact topology of functions. The covariance structure of the matrix-valued Brownian motions $A(t), B(t)$ is as follows. We have

$$A_{ij} = \bar{A}_{ji}, \quad B_{ij} = B_{ji} \quad (10)$$

(B is complex symmetric, not Hermitian) and

$$\mathbf{E}|B_{ij}|^2 = \mathbf{E}A_{ii}A_{jj} = \mathbf{E}A_{ii}\bar{A}_{jj} = \mathbf{E}|A_{ij}|^2 = \langle |O_i|^2, |O_j|^2 \rangle t \quad (11)$$

and all covariances that do not follow from the above are zero.

Remark 5. In the case $G = Z_m$ by (2) we have

$$A(1) = \frac{1}{\sqrt{m+1}}(A' + \zeta I), \quad B(1) = \frac{1}{\sqrt{m+1}}B' \quad (12)$$

Here A' is a version of the GUE with $A'_{i,j} = \bar{A}'_{j,i}$ standard complex normals for $i \neq j$ and mean zero variance $1/2$ i.i.d real normals in the diagonal. (The usual GUE would have variance 1 in the diagonal.) The random variable ζ is standard normal and independent of A' .

B' is an independent symmetric matrix with i.i.d. standard complex normal entries above and below the diagonal and i.i.d. mean zero complex normals with variance $3/2$ in the diagonal (which are independent of everybody else).

Next, we consider the eigenvalues of the operators M_n .

Corollary 6. Consider the operators M_n of Proposition 4. Assume that along some subsequence $\bar{Z}^{n+1} \rightarrow \hat{Z}$ as $n \rightarrow \infty$. Then on this subsequence we have

$$\text{spec}(M_n) \Rightarrow \text{zeros}_\lambda \left[\det \Im(\hat{Z}(Y(1)_{(1,1)} - Y(1)_{(1,2)})) \right].$$

We proceed with the proof of Proposition 4. The proof relies on the noise explosion phenomenon introduced in Section 1.

Proof of Proposition 4. We will assume $\sigma = 1$, the general case may be proved the same way. We first study the evolution without scaling time. From (7)

$$X_k - X_{k-1} = (T_*^{(O \otimes I_2)Q})^{-k} (T_k T_*^{-1} - I)^{(O \otimes I_2)Q} (T_*^{(O \otimes I_2)Q})^k X_{k-1}$$

The coefficient of X_{k-1} above is given by

$$R_k = \begin{bmatrix} Z^k & 0 \\ 0 & Z^{-k} \end{bmatrix} Q^{-1} \begin{bmatrix} 0 & \frac{\lambda}{n}I + \frac{1}{\sqrt{n}}V_k^O \\ 0 & 0 \end{bmatrix} Q \begin{bmatrix} Z^{-k} & 0 \\ 0 & Z^k \end{bmatrix}. \quad (13)$$

Note that the vector $(X_k(\lambda) : \lambda \in \Lambda)$ is a Markov chain in k . In order to prove that it converges to the appropriate SDE we use Proposition 13 of Section 6.

We first rewrite $R_k = \frac{1}{\sqrt{n}}R'_k + \frac{1}{n}R''_k$ where R'_k contains V_k^O and R''_k contains λI from the middle term of R_k . We first focus on the noise term R'_k which expands to

$$R'_k = iS \begin{bmatrix} Z^k & 0 \\ 0 & Z^{-k} \end{bmatrix} \begin{bmatrix} I & -Z \\ -I & \bar{Z} \end{bmatrix} \begin{bmatrix} 0 & V_k^O \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \bar{Z} & Z \\ I & I \end{bmatrix} \begin{bmatrix} Z^{-k} & 0 \\ 0 & Z^k \end{bmatrix} S$$

the three middle factors simplify and we get

$$R'_k = iS \begin{bmatrix} Z^k V_k^O Z^{-k} & Z^k V_k^O Z^k \\ -Z^{-k} V_k^O Z^{-k} & -Z^{-k} V_k^O Z^k \end{bmatrix} S \quad (14)$$

(the diagonal block entries are negative conjugates of each other (note that V_k^O is real) and so are the off-diagonals). For the convergence to the limit, we need to understand the covariance of the partial sums of R_k over k . (This is needed for condition (25) in Proposition 13.)

For this, we may ignore the S factors for the moment, and study

$$A_\ell = \sum_{k=1}^{\ell} Z^k V_k^O Z^{-k}, \quad B_\ell = \sum_{k=1}^{\ell} Z^k V_k^O Z^k \quad (15)$$

The i, j entry of the first term is

$$\sum_{k=1}^{\ell} (z_i \bar{z}_j)^k \sum_r O_{j,r}^T v_{k,r} O_{r,i} = \sum_r \sum_{k=1}^{\ell} v_{k,r} (z_i \bar{z}_j)^k O_{r,j} O_{r,i}$$

Consider the complex covariance of the i, j and i', j' entries in A_ℓ . Since that matrix is Hermitian, we may assume $i \leq j$ and $i' \leq j'$. Since the v 's are independent, the covariance is given by

$$\mathbf{E} A_{i,j} \bar{A}_{i',j'} = \sum_{k=1}^{\ell} (z_i \bar{z}_j \bar{z}_{i'} z_{j'})^k \sum_r O_{r,j} O_{r,i} O_{r,j'} O_{r,i'} = \begin{cases} \ell \langle |O_i|^2, |O_j|^2 \rangle & \text{for } (i, j) = (i', j'), \\ \ell \langle |O_i|^2, |O_{i'}|^2 \rangle & \text{for } (i, i') = (j, j'), \\ \mathcal{O}(1) & \text{otherwise.} \end{cases}$$

This is because we can write $z_i \bar{z}_j \bar{z}_{i'} z_{j'} = e^{i\alpha}$ with $|\alpha| \leq \pi$ and unless $\alpha = 0$ the sum will be $\mathcal{O}(|\alpha|^{-1})$. By the chaoticity condition $\alpha = 0$ can happen only if the conjugated z_i are matched up in pairs to equal non-conjugated z_i , in which case the indices have to be the same. The only options are $i = i', j = j'$ or $i = j, i' = j'$.

This explains that the correlation between an entry A and the entry of B is always $\mathcal{O}(1)$: there will be an odd number of conjugated z 's so such a matching will not occur.

The expectation of the product on the other hand is given by

$$\mathbf{E}A_{i,j}A_{i',j'} = \sum_{k=1}^{\ell} (z_i \bar{z}_j z_{i'} \bar{z}_{j'})^k \sum_r O_{r,j} O_{r,i} O_{r,j'} O_{r,i'} = \begin{cases} \ell \langle O_i^2, O_{i'}^2 \rangle & \text{for } i = j \text{ and } i' = j', \\ \mathcal{O}(1) & \text{otherwise.} \end{cases}$$

indeed, these are the only possible matchings for the conjugated and not-conjugated z 's if $i \leq j$ and $i' \leq j'$. Now

$$\mathbf{E}B_{i,j}B_{i',j'} = \mathcal{O}(1)$$

since no matching can occur in the relevant product $z_i z_j z_{i'} z_{j'}$. Finally, for $i \leq j$ and $i' \leq j'$ (since B is a complex symmetric matrix)

$$\mathbf{E}B_{i,j}\bar{B}_{i',j'} = \begin{cases} \ell \langle O_i^2, O_{j'}^2 \rangle & \text{for } i = i' \text{ and } j = j', \\ \mathcal{O}(1) & \text{otherwise.} \end{cases}.$$

Turning to the drift term R_k'' we first simplify it to get

$$R_k'' = i(\lambda - \lambda_*)S \begin{bmatrix} I & Z^{2k} \\ -Z^{-2k} & -I \end{bmatrix} S. \quad (16)$$

This leads to the estimate

$$i(\lambda - \lambda_*) \sum_{k=1}^{\ell} S \begin{bmatrix} I & Z^{2k} \\ -Z^{-2k} & -I \end{bmatrix} S = i(\lambda - \lambda_*)S \begin{bmatrix} \ell I & \mathcal{O}(1) \\ \mathcal{O}(1) & -\ell I \end{bmatrix} S. \quad (17)$$

Remark 7. From (14) and (16) it is clear that R_k is of the form $\begin{bmatrix} a & b \\ \bar{b} & \bar{a} \end{bmatrix}$ and thus this will hold for $I + R_k$ as well. The product of such matrices will have the same structure, which explains the last assertion of Section 3.

We now turn back to the proof of Proposition 4. For the proof of the convergence, we will use Proposition 13, for which we need to verify the conditions listed there.

- The boundedness of the cubic terms (24) is proved as follows. It is clear that the increments $X_{k+1} - X_k$ are bounded linear functions of λ and the v_i with coefficients given by the entries of X_k . So condition (24) holds as long as X_k is bounded. To ensure that X_k is bounded, we first consider a truncated process in which X_k will stop changing once $\|X_k\| \geq c$. It follows from the proposition that the truncated process converges to the truncated version of the limit, for every $c > 0$. However, as $c \rightarrow \infty$, the solution of the truncated SDE is with high probability equal to the non-truncated one. It follows that the processes converge.

- Conditions (23) and (25) in the truncated process follow from the calculations in the first part of the proof. Note $a^n(t, x)$ is a covariance matrix of dimension $m^2|\Lambda|$. We demonstrate the process of checking its entries. First note that $a^n(t, x)$ is given by

$$a^n(k/n, x) = \mathbf{E} \left[(X_{k+1}^{(n)} - x)_{i,j,\lambda} (X_{k+1}^{(n)} - x)_{i',j',\lambda'} \mid X_k^{(n)} = x \right]$$

and the rows and columns are indexed by $(i, j, \lambda), (i', j', \lambda')$, respectively. Consider the $(i, j, \lambda) = (i', j', \lambda')$ case.

Instead of working directly with a and b , we introduce

$$\hat{Y}_k = \frac{\lambda}{n} \begin{bmatrix} I & Z^{2k} \\ -Z^{-2k} & -I \end{bmatrix} + \frac{1}{\sqrt{n}} \begin{bmatrix} Z^k V_k^O Z^{-k} & Z^k V_k^O Z^k \\ -Z^{-k} V_k^O Z^{-k} & -Z^{-k} V_k^O Z^k \end{bmatrix}$$

and let $\hat{a}^n(t)$ and $\hat{b}^n(t)$ be the n times the second and first moments of the $m^2|\Lambda|$ -vector $\hat{Y}_{\lfloor nt \rfloor}$. Then we have

$$X_k - x = i S \hat{Y}_k S x$$

given $X_{k-1} = x$. This is a linear function of x , and since x is bounded by truncation, it suffices to check (25) for $\hat{a}$ and $\hat{b}$ rather than the original a, b . Similarly, (23) for (the truncated) a, b is implied by

$$|\hat{a}^n(t)| + |\hat{b}^n(t)| \leq c$$

This, in turn, follows from the expression for $\hat{Y}_k$ above (note that Z is diagonal with unit complex numbers).

Returning to (25), we first check the existence of $\hat{b}$ so that

$$\sup_t \left| \int_0^t \hat{b}^n(s) ds - \int_0^t \hat{b}(s) ds \right| \rightarrow 0. \quad (18)$$

This is clear by the following computation, based on (17):

$$\hat{b}(k/n) = \sum_{\ell=1}^k \mathbf{E} \hat{Y}_\ell^{(n)} = \sum_{\ell=1}^k \frac{\lambda}{n} \begin{bmatrix} Z^\ell I Z^{-\ell} & Z^\ell I Z^\ell \\ -Z^{-\ell} I Z^{-\ell} & -Z^{-\ell} I Z^\ell \end{bmatrix} = \lambda \frac{k}{n} \begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix} + \mathcal{O}(n^{-1})$$

where the $\mathcal{O}$ is uniform in k . To check the second moment terms, we first note that since the mean increments are of $\mathcal{O}(1/n)$, it suffices to look at the variance $\tilde{a}(t)$ instead of the second moment $\hat{a}(t)$. Then $\tilde{a}(t)$ is the second moment of

$$\frac{1}{\sqrt{n}} \begin{bmatrix} Z^k V_k^O Z^{-k} & Z^k V_k^O Z^k \\ -Z^{-k} V_k^O Z^{-k} & -Z^{-k} V_k^O Z^k \end{bmatrix}$$

This, by the computations above (starting at equation (14)), converges to the covariance matrix of the corresponding noise term in the SDE.

For the last analyticity statement, we will need a bound of the form

$$\mathbf{E}\|X_k^{(n)}(\lambda)\| \leq f(\lambda, k/n) \quad (19)$$

where $f(\lambda, t)$ is a locally bounded function. Then by Cauchy's theorem it follows that $\max_A \|X_k^{(n)}(\lambda)\|$ is tight for any compact set A , and the claim then follows from Proposition 14. We will follow the squared process instead. We have

$$\begin{aligned} \mathbf{E}(\|X_k\|^2 - \|X_{k-1}\|^2 \mid \mathcal{F}_k) &= \mathbf{E} \operatorname{tr}(\Delta X X_{k-1}^T + \Delta X^T X_{k-1} + \Delta X \Delta X^T) \\ &\leq \frac{c(1 + \lambda^2)}{n} \|X_{k-1}\|^2 \end{aligned}$$

Taking expectations we now see that

$$\mathbf{E}\|X_k\|^2 - \mathbf{E}\|X_{k-1}\|^2 \leq \frac{c(1 + \lambda^2)}{n} \mathbf{E}\|X_{k-1}\|^2$$

rearranging and taking a product for $1, \dots, k$ we immediately get the estimate

$$\mathbf{E}\|X_k\|^2 \leq \left(1 + \frac{c(1 + \lambda^2)}{n}\right)^k \|X_0\|^2 \leq \exp(c(1 + \lambda^2)t) \|X_0\|^2$$

from which the bound (19) follows. $\square$

Proof of Corollary 6. By Skorokhod embedding, we can realize the distributional convergence of the matrix-valued analytic functions $X^{(n)} \Rightarrow Y(1)$ as almost sure convergence. In particular, along our subsequence, we have a.s.

$$\det((\bar{Z}^{n+1}, Z^{n+1})X_n^{(n)}(I, -I)^T) \rightarrow \det((\bar{Z}^*, Z^*)Y_\lambda(1)(I, -I)^T)$$

uniformly on compacts, and the limit is analytic in λ . Note that for $\lambda = 0$ the matrix in the determinant equals

$$\Im(\bar{Z}^*(Y_0(1)_{11} - Y_0(1)_{12})).$$

whose distribution is absolutely continuous with respect to the distribution of the GOE; this follows from the SDE (9). Thus the determinant as a function of λ is not identically zero with probability 1, and the zeros of the subsequence converge to the zeros of the limit almost surely in our realization. The distributional convergence follows. $\square$

5 GOE as a limit

In order to get the GOE limit, we need to take a sequence of boxes of the right dimension. First, some arithmetic conditions have to be satisfied, and by the following lemma they are in most cases.

Lemma 8. *Let λ_* be so that*

$$\frac{2 - \lambda_*}{2 + \lambda_*} \neq \frac{d_j}{d_{j'}} \quad \forall j, j'. \quad (20)$$

Then there exists $r_i \rightarrow 0$ and $n_i \rightarrow \infty$ so that for each r_i the points $\arccos((\lambda_ + r_i d_j)/2)$ are chaotic and for each j , we have $n_i \arccos((\lambda_* + r_i d_j)/2) \rightarrow 0 \pmod{2\pi}$.*

Proof. We will find r_i so that the vector $q = q(r_i) = (\arccos((\lambda_* + r_i d_j)/2))_{j=1, \dots, m}$ has no nonzero integer vector orthogonal to it. It follows that the orbit $\{nq \pmod{2\pi} : n \geq 0\}$ is dense on the m -torus, and we can find the desired convergence to 0.

To find such r_i we show that all but a countable set works. It suffices to show that for any fixed integer vector w , only countably many r_i has $q(r_i) \cdot w = 0$. Note that $r \mapsto q(r) \cdot w$ is an analytic function on the interval $[0, r^*)$ where r^* is the first value so that for some j we have $w_j \neq 0$ and $|\lambda_* + r^* d_j| = 2$. If this happens for a single j , then it follows that the function $r \mapsto q(r) \cdot w$ is not identically zero (since it has a singularity at r^*), so it has countably many zeros.

If this happens for more than one j , then the singularities may cancel out. However, this implies

$$\lambda_* + r^* d_j = -\lambda_* + r^* d_{j'} = 2,$$

which is the reason we banished all such λ_* . □

Proposition 9. *Let G be a real symmetric matrix with distinct eigenvalues d_i , and fix λ_* satisfying (20). Consider*

- *a sequence of $r_\nu \rightarrow r$ so that the critical angles $q_\nu = (\arccos(\lambda_* + r_\nu d_j))_{j=1, \dots, m}$ exist and are linearly independent over the integers,*
- *a sequence n_ν so that for each j we have $(n_\nu + 1)q_{\nu, j} \rightarrow 0 \pmod{2\pi}$ and $n_\nu \text{cha}(q_\nu) \rightarrow \infty$,*
- *a sequence $\sigma_\nu \rightarrow 0$ sufficiently slowly so that we still have $\frac{1}{\sigma_\nu} \max_j (e^{i(n_\nu + 1)q_{\nu, j}} - 1) \rightarrow 0$ and $\sigma_\nu n_\nu \text{cha}(q_\nu) \rightarrow \infty$.*
- *a sequence V_ν of diagonal perturbation matrices where the entries are independent, have mean 0, variance 1 and uniformly bounded third moment.*

Then the regularized transfer matrix evolution for the operator

$$M_\nu = \frac{n_\nu}{\sigma_\nu} \left(rG \times \mathbb{Z}_{n_\nu} + \frac{\sigma_\nu}{\sqrt{n_\nu}} V_\nu - \lambda_* I \right) \quad (21)$$

satisfies the following. For any finite $\Lambda \subset \mathbb{C}$ we have convergence in distribution for the regularized transfer matrices

$$\left(\frac{1}{\sigma_\nu} (X_{[n_\nu t]}^\nu (\lambda^* + \lambda \sigma_\nu / n_\nu) - I), t \in [0, 1], \lambda \in \Lambda \right) \Rightarrow (Y_t(\lambda), t \in [0, 1], \lambda \in \Lambda)$$

where

$$Y = iS^2 \begin{bmatrix} I & 0 \\ 0 & -I \end{bmatrix} \lambda t + iS \begin{bmatrix} A & B \\ -\bar{B} & -\bar{A} \end{bmatrix} S$$

and $S = \lim S_\nu$. Here A, B are matrix-valued Brownian motions with covariance structure given by (10), (11).

Moreover, the eigenvalue process of M_ν converges to the eigenvalues of

$$S_{11} \Im(A(1) - B(1)) S_{11}.$$

Remark 10. In the case when the graph is $\mathbb{Z}_m$, and $S = sI$, the matrix

$$S_{11} \Im(A(1) - B(1)) S_{11}$$

can be written as $(m+1)^{-1/2} s^2 (K + bI)$, where b is a standard normal random variable, and K is a version of the GOE: a real symmetric matrix with independent mean zero real normal entries so that

$$\mathbf{E} K_{ij}^2 = \begin{cases} 1 & i \neq j \\ 5/4 & i = j \end{cases}. \quad (22)$$

Note that for the GOE, the diagonal terms have variance 2. The distribution of the matrix K is not invariant under orthogonal conjugation. The bI term amounts to a random shift of the local eigenvalue process.

Proof. The proof follows very closely the proof of Proposition 4. The only difference is that the frequency of oscillation in the oscillatory sums (15) changes with ν . In particular, for uniform convergence, we need to choose n_ν so that the geometric sums appearing in (15) are $o(\sigma_\nu n_\nu)$. But these sums are bounded by $c/\text{cha}(q_\nu)$, which is still $o(\sigma_\nu n_\nu)$ by our assumptions.

For the convergence of eigenvalues, note that

$$\frac{1}{\sigma_\nu} (\bar{Z}^{n+1}, Z^{n+1}) X_{n_\nu}^{(n)}(I, -I)^T$$

we write the above as

$$\frac{1}{\sigma_\nu}(\bar{Z}^{n+1} - I, Z^{n+1} - I)X_{n_\nu}^{(n)}(I, -I)^T + \frac{1}{\sigma_\nu}(I, I) \cdot (I, -I)^T + \frac{1}{\sigma_\nu}(I, I)(X_{n_\nu}^{(n)} - I)(I, -I)^T$$

The first term converges to 0 since $X_{n_\nu}^{(n)}(I, -I)^T$ is tight and $(Z^{n+1} - I)/\sigma_\nu \rightarrow 0$ by assumption. The second term vanishes, so considering the third term we conclude

$$\frac{1}{\sigma_\nu}(\bar{Z}^{n+1}, Z^{n+1})X_{n_\nu}^{(n)}(I, -I)^T \rightarrow (I, I)Y_\lambda(1)(I, -I)^T$$

all the convergence of the above terms are uniform of compacts. We now take determinants and use the fact that convergence of analytic functions implies the convergence of their zeros, the claim follows. $\square$

Corollary 11. *For almost all $\lambda_* \in [-2, 2]$ there exists a sequence of integer 2-vectors $(m, n) = (m_\nu, n_\nu)$ and $r_\nu \rightarrow 0, \sigma_\nu, \gamma_\nu$ so that the eigenvalue process of*

$$\gamma[(r\mathbb{Z}_m) \times \mathbb{Z}_n + \sigma V - \lambda_* I]$$

as $\nu \rightarrow \infty$ converges locally to the Sine₁ process. Here the noise matrices V_ν are diagonal with independent entries having arbitrary fixed mean 0 distribution with uniformly bounded third moment.

Proof. As $n \rightarrow \infty$, by the methods of Erdős et al. (2009b) the bulk eigenvalue process of the matrices $K + bI$ (see Remark 10) converges locally to the Sine₁ process. The proof of this statement will be expanded in Lemma 16 in the Appendix.

We now consider a λ_* so that

$$\frac{2 - \lambda_*}{2 + \lambda_*} \neq \frac{\cos(q_1\pi)}{\cos(q_2\pi)}$$

for any rational q_1, q_2 . Consider a base graph $G = \mathbb{Z}_m$. Then by Proposition 9 and Lemma 8 we can find sequences $n_\nu, r_\nu \rightarrow 0, \sigma_\nu$ so that the eigenvalues of (21) M_n converge to those of $s^2(K + bI)$. After we have done this for a sequence of base graphs with $m \rightarrow \infty$, we can find a diagonal sequence along which, after the appropriate scaling, the desired convergence holds. $\square$

Remark 12 (Higher dimensional boxes). When $G = \mathbb{Z}_{m_1} \times \dots \times \mathbb{Z}_{m_d}$ then the eigenvectors of G are of the form

$$O_{j_1, \dots, j_d}(k_1, \dots, k_d) = \prod_{\ell=1}^d \sqrt{\frac{2}{m_\ell + 1}} \sin\left(\frac{\pi j_\ell k_\ell}{m_\ell + 1}\right)$$

and the corresponding eigenvalues are $2 \sum_{\ell=1}^d \cos\left(\frac{j_\ell \pi}{1+m_\ell}\right)$. One can also calculate that

$$\left(\prod_{\ell=1}^d (m_\ell + 1)\right) \langle |O_{\underline{i}}|^2, |O_{\underline{j}}|^2 \rangle = \prod_{\ell=1}^d \left(1 + \frac{1}{2} 1(i_\ell = j_\ell)\right).$$

If the eigenvalues are distinct and λ^* satisfies (20) then Proposition 9 will still apply. However, the limiting symmetric real Gaussian matrix $\Im(A(1) - B(1))$ will have a more complicated covariance structure for $d > 1$ then the one described in Remark 10 and the current results are not strong enough to imply that the bulk scaling limit will be the Sine₁ process.

One can check that the random matrix still contains an order 1 component of a GOE matrix which means that the local relaxation flow argument of Erdős et al. (2009b) will go through. The problem is caused by the fact that for the ‘other’ component the strong local semicircle result is not available.

6 Appendix

6.1 SDE limit of Markov chains

The following is the main tool for proving convergence in the oscillatory setting. It is Proposition 23 in Valkó and Virág (2009), and is based on Theorem 7.4.1 of Ethier and Kurtz (1986).

Proposition 13. *Fix $T > 0$, and for each $n \geq 1$ consider a Markov chain*

$$(X_\ell^n \in \mathbb{R}^d, \ell = 1 \dots \lfloor nT \rfloor).$$

Let $Y_\ell^n(x)$ be distributed as the increment $X_{\ell+1}^n - x$ given $X_\ell^n = x$. We define

$$b^n(t, x) = nE[Y_{\lfloor nt \rfloor}^n(x)], \quad a^n(t, x) = nE[Y_{\lfloor nt \rfloor}^n(x)Y_{\lfloor nt \rfloor}^n(x)^T].$$

Suppose that as $n \rightarrow \infty$ we have

$$|a^n(t, x) - a^n(t, y)| + |b^n(t, x) - b^n(t, y)| \leq c|x - y| + o(1) \quad (23)$$

$$\sup_{x, \ell} E[|Y_\ell^n(x)|^3] \leq cn^{-3/2}, \quad (24)$$

and that there are functions a, b from $\mathbb{R} \times [0, T]$ to $\mathbb{R}^{d^2}, \mathbb{R}^d$ respectively with bounded first and second derivatives so that

$$\sup_{x, t} \left| \int_0^t a^n(s, x) ds - \int_0^t a(s, x) ds \right| + \sup_{x, t} \left| \int_0^t b^n(s, x) ds - \int_0^t b(s, x) ds \right| \rightarrow 0. \quad (25)$$

Assume also that the initial conditions converge weakly:

$$X_0^n \Rightarrow X_0.$$

Then $(X_{[nt]}^n, 0 \leq t \leq T)$ converges in law to the unique solution of the SDE

$$dX = b dt + g dB, \quad X(0) = X_0$$

where B is a d -dimensional standard Brownian motion and g is any C^2 function satisfying $gg^T = a$.

We need a result that strengthens the convergence of random analytic functions.

Proposition 14. *Let f_n be a sequence of random analytic functions on a domain D so that $\max_A f_n$ is tight for every compact $A \subset D$ and $f_n \Rightarrow f$ in the sense of finite dimensional distributions. Then f has a unique analytic version and $f_n \Rightarrow f$ in distribution with respect to local-uniform convergence.*

Proof. Pick a countable dense set of points $D' \subset D$. We can first find a subsequence so that the joint distribution of $\max_A |f_n|$ and $f_n(z), z \in D'$ converges. By the Skorokhod's embedding theorem we can realize the sequence $(f_n(z), z \in D')$ on a single probability space Ω so that almost surely $f_n(z) \rightarrow f(z)$ for all $z \in D'$ and also $\max_A |f_n|$ converges (and so it is bounded) for $A \subset \mathcal{A}$. By continuity, we can define the f_n as analytic functions on D for the probability space Ω . Then the f_n have at least one analytic limit in the sense of uniform-on-compacts convergence by Montel's theorem. This limit must agree with f on all points $z \in D'$, so it is unique.

By the above argument any sequence has a further subsequence that converges locally uniformly to some analytic f in distribution. But the distribution of f is determined by its finite dimensional distributions, so the limit is unique. $\square$

Corollary 15. *Let f_n be random analytic functions on D so that $\mathbf{E}h(|f_n(z)|) \leq g(z)$ for some and increasing unbounded function h and locally bounded function g . Assume that $f_n(z) \Rightarrow f(z)$ in the sense of finite dimensional distributions. Then f has a unique analytic version and $f_n \Rightarrow f$ in distribution with respect to local-uniform convergence.*

Proof. By Cauchy's theorem $\max_A |f_n|$ is tight for every compact set A , and the claim follows from the Proposition. $\square$

6.2 The point process limit of the modified GOE

Consider the $n \times n$ symmetric random matrix $H_n = n^{-1/2}(K + bI)$ where b is a standard normal random variable, and K is the version of the GOE considered in (22). We will show that the eigenvalue process of H_n in the bulk converges locally to the Sine₁ process, the bulk scaling limit of the GOE eigenvalue process. Bulk scaling means that we consider $\rho(\lambda)\sqrt{n}(H_n - \lambda I)$ where $|\lambda| < 2$ and $\rho(\lambda) = \frac{1}{2\pi}\sqrt{4 - \lambda^2} 1_{|\lambda| \leq 2}$ is the semicircle density.

Lemma 16. *For any $|\lambda| < 2$ and compactly supported continuous test function $\Theta : \mathbb{R}^k \rightarrow \mathbb{R}$ we have*

$$\int_{\mathbb{R}^n} d\alpha_1 \dots d\alpha_n \Theta(\alpha_1, \dots, \alpha_k) \rho(\lambda)^{-k} (p_{H,n}^{(k)} - p_{GOE,n}^{(k)}) \left(\lambda + \frac{\alpha_1}{n\rho(\lambda)}, \dots, \lambda + \frac{\alpha_k}{n\rho(\lambda)} \right) \rightarrow 0. \quad (26)$$

Here $p_{H,n}^{(k)}$ is the k -point intensity function of the eigenvalues of H_n . The function $p_{GOE,n}^{(k)}$ is the same for the $n \times n$ GOE matrix with variances $2/n$ and $1/n$ on the diagonal and elsewhere, respectively.

Proof. Since the proof works exactly the same way for any $|\lambda| < 2$, we will assume $\lambda = 0$.

Our proof relies on the local relaxation flow arguments of Erdős et al. (2009b) and the strong local semicircle law proved in Erdős et al. (2010). The argument in a nutshell is the following: if we have a real symmetric random matrix whose eigenvalues are well-approximated by the semicircle law locally then by adding a small constant times an independent GOE matrix (i.e. by running Dyson's Brownian motion for a small time with our matrix as the initial condition) the scaled eigenvalue process of the resulting matrix will be close to the Sine₁ process.

We will use Theorem 2.3 of Erdős et al. (2010) (see also the comments after the theorem) which provides a powerful quantitative version of the previous argument. Since we can write $\frac{1}{\sqrt{n}}K = K_1 + K_2$ where K_1 is $1/2$ times a GOE and K_2 is an independent symmetric random matrix satisfying the conditions of the theorem then for any $q > 0$ we have

$$\begin{aligned} & \left| \int_{-\frac{q}{\sqrt{n}}}^{\frac{q}{\sqrt{n}}} \frac{d\mu\sqrt{n}}{2q} \int_{\mathbb{R}^k} d\alpha_1 \dots d\alpha_n \Theta(\alpha_1, \dots, \alpha_k) \rho(0)^{-k} \right. \\ & \quad \left. \times (p_{K,n}^{(k)} - p_{GOE,n}^{(k)}) \left(\mu + \frac{\alpha_1}{n\rho(0)}, \dots, \mu + \frac{\alpha_k}{n\rho(0)} \right) \right| \leq Cn^{-1/4+\varepsilon} (q^{-1} + q^{-1/2}). \end{aligned} \quad (27)$$

(We apply the theorem with $E = 0, b = q/\sqrt{n}, t = 1/2, \varepsilon' \ll 1$ and $\delta = 1 - \varepsilon'$.) Since the spectrum of H_n can be obtained by a random shift of the spectrum of $n^{-1/2}K$ we have

$$p_{H,n}^{(k)} \left(\frac{\alpha_1}{n\rho(0)}, \dots, \frac{\alpha_k}{n\rho(0)} \right) = \int_{-\infty}^{\infty} \frac{db}{\sqrt{2\pi n^{-1}}} e^{-\frac{b^2 n}{2}} p_{K,n}^{(k)} \left(b + \frac{\alpha_1}{n\rho(0)}, \dots, b + \frac{\alpha_k}{n\rho(0)} \right). \quad (28)$$

By Fubini for any nonnegative function F we have

$$\int_{-\infty}^{\infty} \frac{db}{\sqrt{2\pi n^{-1}}} e^{-\frac{b^2 n}{2}} F(b) = \int_0^{\infty} e^{-\frac{q^2}{2}} \sqrt{\frac{2}{\pi}} q^2 \int_{-\frac{q}{\sqrt{n}}}^{\frac{q}{\sqrt{n}}} \frac{db \sqrt{n}}{2q} F(b) dq.$$

Using this with (27) and (28) we get the upper bound

$$|\text{left hand side of (26)}| \leq \int_0^{\infty} e^{-\frac{q^2}{2}} \sqrt{\frac{2}{\pi}} q^2 C n^{-1/4+\varepsilon} (q^{-1} + q^{-1/2}) \leq C' n^{-1/4+\varepsilon}$$

which shows that the limit of the eigenvalue process of H_n is the same as the limit of *GOE* eigenvalues multiplied by n and shifted by an independent Gaussian b_n of variance n . We first condition on the sequence of these independent Gaussians, so the GOE eigenvalues are now centered at $-b_n$, which, on the semicircle scale, converges to 0. In the classical literature, convergence is usually proved around a fixed window at c , $|c| < 2$ (on the semicircle scale). The moving window case is rigorously proved in Valkó and Virág (2009). The lemma follows after removing the conditioning. $\square$

Acknowledgements. We thank Eugene Kritchanski, Alex Bloemendal, and László Erdős for useful discussions, and Hermann Schulz-Baldes for references. The second author thanks the hospitality of the Banff International Research Station, where part of this work was conceived. This research is supported by the NSF Grant DMS-09-05820 (Valkó) and the NSERC discovery accelerator grant program and the Canada Research Chair program (Virág).

References

- B. Altshuler and B. I. Shklovski (1986). Repulsion of energy levels and conductivity of metal samples. *Sov. Phys. JETP*, 64:127–135.
- P. W. Anderson (1958). Absence of diffusion in certain random lattices. *Phys. Rev.*, 109: 1492–1505.
- S. Bachmann and W. De Roeck (2010). From the Anderson model on a strip to the DMPK equation and random matrix theory. *J. Stat. Phys.* 139, no. 4, 541564.
- O. Bohigas, M.-J. Giannoni, and C. Schmit (1984). Characterization of chaotic quantum spectra and universality of level fluctuation laws. *Phys. Rev. Lett.*, 52(1):1–4.
- K. B. Efetov. *Supersymmetry in disorder and chaos*. Cambridge UP, 1997.

- L. Erdős, B. Schlein, and H.-T. Yau. Universality of random matrices and local relaxation flow, 2009a. [arXiv.org:0907.5605](https://arxiv.org/abs/0907.5605).
- L. Erdős, B. Schlein, H.-T. Yau, and J. Yin. The local relaxation flow approach to universality of the local statistics for random matrices, 2009b. [arXiv.org:0911.3687](https://arxiv.org/abs/0911.3687).
- L. Erdős, H.-T. Yau, and J. Yin. Rigidity of Eigenvalues of Generalized Wigner Matrices, 2010. [arXiv.org:1007.4652](https://arxiv.org/abs/1007.4652).
- S. N. Ethier and T. G. Kurtz. *Markov processes*. John Wiley & Sons Inc., New York, 1986.
- E. Kritchevski, B. Valkó, and B. Virág. The scaling limit of the critical one-dimensional random Schrödinger operator, 2011. [arXiv.org:1107.3058](https://arxiv.org/abs/1107.3058).
- M. L. Mehta. *Random matrices*. Elsevier/Academic Press, Amsterdam, third edition, 2004.
- R.A. Römer, R.A. and H. Schulz-Baldes (2010). The random phase property and the Lyapunov spectrum for disordered multi-channel systems. *Journal of Statistical Physics*, 140. no. 1. 122–153.
- H. Schulz-Baldes (2004). Perturbation theory for Lyapunov exponents of an Anderson model on a strip, *GAFSA*. 14, 1089–1117.
- T. Tao and V. Vu. Random matrices: Universality of local eigenvalue statistics, 2009. [arXiv.org:0906.0510](https://arxiv.org/abs/0906.0510).
- B. Valkó and B. Virág (2009). Continuum limits of random matrices and the Brownian carousel. *Inventiones Math.*, 177:463–508.
- E. P. Wigner (1957). *Gatlinberg Conference on Neutron Physics, Oak Ridge National Laboratory Report*, ORNL 2309:59.